

## KCTD4 蛋白抗体

产品货号： mIR16935

英文名称： KCTD4

中文名称： KCTD4 蛋白抗体

别名： bA321C24.3; BTB/POZ domain containing protein KCTD4; KCTD 4; OTTHUMP00000018343;  
Potassium channel tetramerisation domain containing 4.

研究领域： 肿瘤 细胞生物 信号转导 转录调节因子

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Chicken, Dog, Pig, Cow, Horse, Rabbit, Sheep,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需  
做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量 : 30kDa

细胞定位 : 细胞膜

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human KCTD4:161-259/259

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

**PubMed :** PubMed

**产品介绍 :** KCTD4 contains a potassium channel tetramerisation domain. The N terminal, cytoplasmic tetramerisation domain (T1) of voltage-gated potassium channels encodes molecular determinants for subfamily specific assembly of alpha subunits into functional tetrameric channels. The specific function of KCTD4 is unknown.

**Similarity:**

Contains 1 BTB (POZ) domain.

**SWISS:**

Q8WVF5

**Gene ID:**

386618

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.