

补体 C1q 肿瘤坏死因子相关蛋白 6 抗体

产品货号： mlR14104

英文名称： CTRP6

中文名称： 补体 C1q 肿瘤坏死因子相关蛋白 6 抗体

别名： C1q and tumor necrosis factor related protein 6; C1QT6_HUMAN; C1QTNF6; Complement C1q tumor necrosis factor related protein 6; Complement C1q tumor necrosis factor-related protein 6; CTRP 6; CTRP6; OTTHUMP00000197772; OTTHUMP00000197774; PRO1151; UNQ581; ZACRP6; ZACRP6.

研究领域： 细胞生物 免疫学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat,

产品应用： WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500

（石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 42kDa

细胞定位： 细胞浆

性状： Lyophilized or Liquid

浓度： 1mg/ml

免疫原： KLH conjugated synthetic peptide derived from human CTRP6:151-259/259

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PubMed : PubMed

产品介绍 : CTRP6 is a 259 amino acid secreted glycoprotein consisting of an N-terminal signal peptide sequence followed by a collagen-like domain and a C-terminal globular domain. Predominantly expressed in eye and placenta, CTRP6 is also upregulated in obese mice. Along with CTRP1, CTRP2, CTRP3 and CTRP5, CTRP6 circulates in the blood and is thought to function as an endocrine hormone. There are three isoforms of CTRP6 that are produced as a result of alternative splicing events.

Subcellular Location:

Secreted.

Similarity:

Contains 1 C1q domain.

Contains 1 collagen-like domain.

SWISS:

Q9BXI9

Gene ID:

114904

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

产品图片

