



小鼠抗拓普西异构酶 I 单克隆抗体

产品货号 : mlR51396

英文名称 : Topoisomerase I

中文名称 : 小鼠抗拓普西异构酶 I 单克隆抗体

别 名 : DNA topoisomerase 1; DNA topoisomerase I; NUP98 fusion gene; TOP 1; TOP I; TOP1; TOP1_HUMAN; TOPI; Topoisomerase 1; Topoisomerase1; Topoisomerasel; Type I DNA topoisomerase.

研究领域 : 细胞生物 染色质和核信号

抗体来源 : Mouse

克隆类型 : Monoclonal

克 隆 号 : 3D7

交叉反应 : Human, Mouse, Rat,

产品应用 : WB=1:500-2000

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量 : 91kDa

细胞定位 : 细胞核

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human Topoisomerase I:



亚 型 : IgG1

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed

产品介绍 : This gene encodes a DNA topoisomerase, an enzyme that controls and alters the topologic states of DNA during transcription. This enzyme catalyzes the transient breaking and rejoining of a single strand of DNA which allows the strands to pass through one another, thus altering the topology of DNA. This gene is localized to chromosome 20 and has pseudogenes which reside on chromosomes 1 and 22. [provided by RefSeq, Jul 2008]

Function:

The reaction catalyzed by topoisomerases leads to the conversion of one topological isomer of DNA to another.

Subcellular Location:

Nucleus, nucleolus. Nucleus, nucleoplasm. Diffuse nuclear localization with some enrichment in nucleoli. On CPT treatment, cleared from nucleoli into nucleoplasm. Sumolyated forms found in both nucleoplasm and nucleoli.

Post-translational modifications:

Sumoylated. Lys-117 is the main site of sumoylation. Sumoylation plays a role in partitioning TOP1 between nucleoli and nucleoplasm. Levels are dramatically increased on camptothecin (CPT) treatment.

**DISEASE:**

Note=A chromosomal aberration involving TOP1 is found in a form of therapy-related myelodysplastic syndrome.
Translocation t(11;20)(p15;q11) with NUP98.

Similarity:

Belongs to the eukaryotic type I topoisomerase family.

SWISS:

P11387

Gene ID:

7150

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

产品图片

