

跨膜 γ 羧基蛋白 2 抗体

产品货号： mIR19426

英文名称： PRRG2

中文名称： 跨膜 γ 羧基蛋白 2 抗体

别名： PRGP2; Proline rich gamma carboxyglutamic acid protein 2; Proline rich Gla (G carboxyglutamic acid) polypeptide 2; Proline rich Gla protein 2; TMG2; Transmembrane gamma carboxyglutamic acid protein 2.

研究领域： 细胞生物 免疫学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Dog, Pig,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 17kDa

细胞定位： 细胞膜

性状： Lyophilized or Liquid

浓度： 1mg/ml

免疫原 : KLH conjugated synthetic peptide derived from human PRRG2:21-120/202 <Extracellular>

亚型 : IgG

纯化方法 : affinity purified by Protein A

储存液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed

产品介绍 : PRRG2 interacts with NEDD4 and is highly expressed in the thyroid.

Subunit:

Interacts with NEDD4.

Subcellular Location:

Cell Membrane; Single-pass type I membrane protein.

Similarity:

Contains 1 Gla (gamma-carboxy-glutamate) domain.

SWISS:

O14669



Gene ID:

5639

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.