



## 乙酰化和磷酸化组蛋白 H3 抗体

产品货号 : mlR3748

英文名称 : Histone H3 (acetyl K9, phospho S10)

中文名称 : 乙酰化和磷酸化组蛋白 H3 抗体

别 名 : Histone H3 (phospho S10, acetyl K9); Acetyl and phospho-Histone H3 (Ac-K9/p-Ser10); Histone H3 (Ac-K9/p-Ser10); Histone H3 (Acetyl-K9/phospho-Ser10); Histone H3 (Ac-Lys9/p-Ser10); Histone H3 (Acetyl-Lys9/phospho-Ser10); H3 histone family member E pseudogene; H3F3; HIST3H3; Histone H3 3 pseudogene; H31\_TETTH; Histone H3; H3S; Histone H3-I/H3-II; Major histone H3; H3F.

产品类型 : 磷酸化抗体 乙酰化抗体

研究领域 : 肿瘤 细胞生物 免疫学 发育生物学 染色质和核信号 信号转导 转录调节因子

抗体来源 : Rabbit

克隆类型 : Polyclonal

交叉反应 : Human, Mouse, Rat, Pig, Cow, Rabbit, Fruit Fly,

产品应用 : WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 (石蜡切片  
需做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量 : 15kDa

细胞定位 : 细胞核

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml



**免 疫 原 :** KLH conjugated synthetic peptide derived from human Histone H3 around the site of Acetyl-Lys9 and phospho-Ser10:AR(Ac-K)(p-S)TG

**亚 型 :** IgG

**纯化方法 :** affinity purified by Protein A

**储 存 液 :** 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

**保存条件 :** Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

**PubMed :** PubMed

**产品介绍 :** Modulation of the chromatin structure plays an important role in the regulation of transcription in eukaryotes. The nucleosome, made up of four core histone proteins (H2A, H2B, H3 and H4), is the primary building block of chromatin. The N-terminal tail of core histones undergoes different posttranslational modifications including acetylation, phosphorylation and methylation. These modifications occur in response to cell signal stimuli and have a direct effect on gene expression. In most species, the histone H2B is primarily acetylated at lysines 5, 12, 15 and 20. Histone H3 is primarily acetylated at lysines 9, 14, 18 and 23. Acetylation at lysine 9 appears to have a dominant role in histone deposition and chromatin assembly in some organisms. Phosphorylation at Ser10 of histone H3 is tightly correlated with chromosome condensation during both mitosis and meiosis.

**Function:**

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. H3 is deposited into chromatin exclusively through a DNA replication-coupled pathway that can be associated with either DNA duplication or DNA repair synthesis during meiotic homologous recombination.

**Subunit:**

The nucleosome is a histone octamer containing two molecules each of H2A, H2B, H3 and H4 assembled in one H3-H4 heterotetramer and two H2A-H2B heterodimers. The octamer wraps approximately 147 bp of DNA. Interacts with GCN5, whereby H3S10ph increases histone-protein interactions. Interacts with PDD1 and PDD3.

**Subcellular Location:**

Nucleus. Chromosome. Note=Localizes to both the large, transcriptionally active, somatic macronucleus (MAC) and the small, transcriptionally inert, germ line micronucleus (MIC).

**Post-translational modifications:**

Phosphorylated to form H3S10ph. H3S10ph promotes subsequent H3K14ac formation by GCN5. H3S10ph is only found in the mitotically dividing MIC, but not in the a mitotically dividing MAC. H3S10ph is correlated with chromosome condensation during mitotic or meiotic micronuclear divisions.

Acetylation of histone H3 leads to transcriptional activation. H3K14ac formation by GCN5 is promoted by H3S10ph. H3K9acK14ac is the preferred acetylated form of newly synthesized H3. Acetylation occurs almost exclusively in the MAC.

Methylated to form H3K4me. H3K4me is only found in the transcriptionally active MAC. Methylated to form H3K9me in developing MACs during conjugation, when genome-wide DNA elimination occurs. At this stage, H3K9me specifically occurs on DNA sequences being eliminated (IES), probably targeted by small scan RNAs (scRNAs) bound to IES, and is required for efficient IES elimination. H3K9me is required for the interaction with the chromodomains of PDD1 and PDD3.

The full-length protein H3S (slow migrating) is converted to H3F (fast migrating) by proteolytic removal of the first 6 residues. H3F is unique to MIC, and processing seems to occur regularly each generation at a specific point in the cell cycle.

**Similarity:**

Belongs to the histone H3 family.



**SWISS:**

P68431

**Gene ID:**

8290

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.