

酪蛋白激酶 II β 亚基抗体

产品货号 : mlR18453

英文名称 : LY6G5B/C6orf19

中文名称 : 酪蛋白激酶 II β 亚基抗体

别名 : C6orf19; Casein kinase II subunit beta; Chromosome 6 open reading frame 19; CK II beta; CK2N; G5A; G5b; LY65B_HUMAN; LY6G5B; Lymphocyte antigen 6 complex locus protein G5b; Lymphocyte antigen 6 complex, locus G5B; Phosvitin.

研究领域 : 细胞生物 免疫学 激酶和磷酸酶

抗体来源 : Rabbit

克隆类型 : Polyclonal

交叉反应 : Human,

产品应用 : ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 (石蜡切片需做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量 : 21kDa

细胞定位 : 分泌型蛋白

性状 : Lyophilized or Liquid

浓度 : 1mg/ml

免疫原 : KLH conjugated synthetic peptide derived from human LY6G5B/C6orf19:121-201/201

亚型 : IgG

纯化方法 : affinity purified by Protein A

储存液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed

产品介绍 : LY6G5B belongs to a cluster of leukocyte antigen-6 (LY6) genes located in the major histocompatibility complex (MHC) class III region on chromosome 6. Members of the LY6 superfamily typically contain 70 to 80 amino acids, including 8 to 10 cysteines. Most LY6 proteins are attached to the cell surface by a glycosylphosphatidylinositol (GPI) anchor that is directly involved in signal transduction (Mallya et al., 2002 [PubMed 12079290]).[supplied by OMIM, Mar 2008]

Subcellular Location:

Secreted.

Post-translational modifications:

N-glycosylated.

Similarity:

Contains 1 UPAR/Ly6 domain.

SWISS:



Q8NDX9

Gene ID:

58496

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.