

## 转录因子 ER81 蛋白抗体

产品货号： mlR11745

英文名称： ER81

中文名称： 转录因子 ER81 蛋白抗体

别名： DKFZp781L0674; ER81; ER81 protein; ETS translocation variant 1; ets variant gene 1; Ets-related protein 81; Etsrp81; ETV 1; ETV1; ETV1\_HUMAN; MGC104699; MGC120533; MGC120534.

研究领域： 细胞生物 神经生物学 干细胞 表观遗传学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Dog, Pig, Cow, Horse, Sheep,

产品应用： WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500

(石蜡切片需做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 55kDa

细胞定位： 细胞核

性状： Lyophilized or Liquid

浓度： 1mg/ml

免疫原： KLH conjugated synthetic peptide derived from human ER81:21-120/477

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PubMed : PubMed

产品介绍 : Several members of the Ets gene family encode sequence-specific DNA binding proteins that recognize DNA sequences with a centrally located 5'-GGAA-3' element. All of the Ets proteins recognize the same central core sequence but each protein interacts with unique sequences that flank this core. PEA3 binds the motif 5'-AGGAAG-3', while ER81 (also designated ETV1) binds the motif 5'-CGGAA/T-3'. PEA3 is expressed at readily detectable levels in cells of epithelial and fibroblastic origin. Unlike other members of the Ets family, including Ets-1 and Ets-2, PEA3 is not expressed in hematopoietic cells. ER81 is highly expressed in brain, testis, lung and heart. ER81 is also moderately expressed in spleen, pancreas, colon and small intestine. During development, ER81, PEA3 and ERM display unique expression patterns which suggest these transcriptional factors play an important role in organogenesis. ERK-1 activates ER81 transcriptional activity, while MAPKAP kinase 2 inhibits ER81.

**Function:**

Transcriptional activator that binds to DNA sequences containing the consensus pentanucleotide 5'-CGGA[AT]-3'.

**Subcellular Location:**

Nucleus.

**Tissue Specificity:**

Very highly expressed in brain, highly expressed in testis, lung and heart, moderately in spleen, small intestine, pancreas and colon, weakly in liver, prostate and thymus, very weakly in skeletal muscle, kidney and ovary and not in placenta and peripheral blood leukocytes.

**Post-translational modifications:**

Sumoylated.

**DISEASE:**

Defects in ETV1 are a cause of Ewing sarcoma (ES) [MIM:612219]. A highly malignant, metastatic, primitive small round cell tumor of bone and soft tissue that affects children and adolescents. It belongs to the Ewing sarcoma family of tumors, a group of morphologically heterogeneous neoplasms that share the same cytogenetic features. They are considered neural tumors derived from cells of the neural crest. Ewing sarcoma represents the less differentiated form of the tumors. Note=A chromosomal aberration involving ETV1 is found in patients with Erwing sarcoma. Translocation t(7;22)(p22;q12) with EWSR1.

**Similarity:**

Belongs to the ETS family.

Contains 1 ETS DNA-binding domain.

**SWISS:**

P50549

**Gene ID:**

2115

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

产品图片

