

胆碱磷酸酯酶 4 抗体

产品货号： mIR12718

英文名称： PLD4

中文名称： 胆碱磷酸酯酶 4 抗体

别 名： C14orf175; Choline phosphatase 4; EC 3.1.4.4; Phosphatidylcholine hydrolyzing phospholipase D4; Phosphatidylcholine-hydrolyzing phospholipase D4; Phospholipase D family member 4; Phospholipase D4; PLD 4; PLD4; PLD4_HUMAN.

研究领域： 细胞生物 免疫学 信号转导

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Cow, Sheep,

产品应用： IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量： 56kDa

细胞定位： 细胞膜

性 状： Lyophilized or Liquid

浓 度： 1mg/ml

免 疫 原： KLH conjugated synthetic peptide derived from human PLD4:201-300/506

亚 型： IgG

纯化方法： affinity purified by Protein A

储 存 液： 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件： Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed： PubMed

产品介绍： PLD4 (Phospholipase D Family Member 4) is a Protein Coding gene. Among its related pathways are Metabolism and Immune System. GO annotations related to this gene include phosphatidylinositol phospholipase C activity and N-acylphosphatidylethanolamine-specific phospholipase D activity. An important paralog of this gene is PLD3.

Subcellular Location:

Membrane.

Similarity:

Belongs to the phospholipase D family.

Contains 2 PLD phosphodiesterase domains.

SWISS:

Q96BZ4

Gene ID:

122618

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.