

胶质瘤发病相关蛋白 1 样蛋白 2 抗体

产品货号： mIR18920

英文名称： GLIPR1L2

中文名称： 胶质瘤发病相关蛋白 1 样蛋白 2 抗体

别 名： GLI pathogenesis related 1 like 2; GLIPR1-like protein 2; GLIPR1L 2; Glipr1l2; GRPL2_HUMAN.

研究领域： 细胞生物 免疫学 神经生物学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量： 40, 25kDa

细胞定位： 细胞膜

性 状： Lyophilized or Liquid

浓 度： 1mg/ml

免 疫 原： KLH conjugated synthetic peptide derived from human GLIPR1L2:101-200/344

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed

产品介绍 : This gene encodes a member of the cysteine-rich secretory protein, antigen 5, and pathogenesis-related 1 superfamily. Members of this family have roles in a variety of processes, including cancer and immune defense. This gene is located in a cluster with two related genes on chromosome 12. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2012]

Subcellular Location:

Membrane.

Tissue Specificity:

Highly expressed in testis. Detected in prostate, kidney, bladder, lung and bone marrow.

Similarity:

Belongs to the CRISP family.

SWISS:

Q4G1C9

Gene ID:

144321

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.