

微管相关蛋白 NICN1 抗体

产品货号： mIR19239

英文名称： NICN1

中文名称： 微管相关蛋白 NICN1 抗体

别名： Nicolin 1; NPCEDRG; p24; PGs5; Tubulin polyglutamylase complex subunit 5.

研究领域： 细胞生物 免疫学 信号转导 细胞骨架

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Pig, Cow, Horse, Rabbit,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 24kDa

细胞定位： 细胞核

性状： Lyophilized or Liquid

浓度： 1mg/ml

免疫原： KLH conjugated synthetic peptide derived from human NICN1:121-213/213

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed

产品介绍 : This protein encoded by this gene localizes to the nucleus and is expressed in numerous tissues including brain, testis, liver, and kidney. This refseq contains genomic sequence in its 3' UTR which is not supported by experimental evidence. Computer predictions indicate that this region of the 3' UTR contains hairpin-forming self-complementary sequence which is possibly excised after transcription. This gene has a pseudogene on chromosome X. [provided by RefSeq, Jul 2008]

Subcellular Location:

Nuclear

Tissue Specificity:

The function of NICN1 remains unknown. It is localized to the nucleus and is expressed in numerous tissues including brain, testis, liver, and kidney.

SWISS:

Q9BSH3

Gene ID:

84276

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.