

## 雌激素调节基因 1 抗体

产品货号： mIR19920

英文名称： CUZD1

中文名称： 雌激素调节基因 1 抗体

别名： CUB and zona pellucida-like domain-containing protein 1: CUB and ZP domain-containing protein 1:  
CUZD1: ERG-1: Transmembrane protein UO-44: UNQ224/PRO257: UO-44.

研究领域： 肿瘤 免疫学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需  
做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 68kDa

细胞定位： 细胞浆 细胞膜 细胞外基质

性状： Lyophilized or Liquid

浓度： 1mg/ml

**免疫原** : KLH conjugated synthetic peptide derived from human SERF1:61-160/607 <Extracellular>

**亚型** : IgG

**纯化方法** : affinity purified by Protein A

**储存液** : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

**保存条件** : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

**PubMed** : PubMed

**产品介绍** : CUZD1 is a 607 amino acid single-pass type I membrane protein that contains two CUB domains and a ZP (zona-pellucida) domain. Expressed in pancreas and epithelium of ovary, CUZD1 is found at high levels in ovarian tumors where it is thought to inhibit cell attachment and proliferation. CUZD1 is suggested to play a role in cell motility, cell-cell interactions and/or interactions with the extracellular matrices. Existing as three alternatively spliced isoforms, CUZD1 is encoded by a gene located on human chromosome 10. Making up 4.5% of the human genome, chromosome 10 encodes roughly 800 genes including PTEN, a tumor suppressor gene that has been linked to the development of Cowden syndrome. The chromosome 10 encoded gene ERCC6 is important for DNA repair and is linked to Cockayne syndrome which is characterized by extreme photosensitivity and premature aging.

**Function:**

CUZD1 antiserum inhibits cell attachment and proliferation of ovarian cancer cells so may be involved in these processes. May also play a role in the uterus during late pregnancy and/or in trypsin activation in pancreatic acinar cells.

**Subcellular Location:**

Cytoplasmic vesicle > secretory vesicle membrane. Associated with zymogen granule membranes.

**Tissue Specificity:**

Detected in pancreas and epithelium of ovary. Expressed at higher levels in ovarian tumors than in normal tissue.

**Similarity:**

Contains 2 CUB domains.

Contains 1 ZP domain.

**SWISS:**

Q86UP6

**Gene ID:**

50624

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.